



Phormone Chemicals

The name you can always trust

Mfrs: Phormone Traps, Lures, Yellow sticky traps

Chilo partellus (Spotted stalk borer)

Host range – Sorghum, Maize, Bulrush millet, Sugarcane, Rice



Nature of Damage:

In young plants the shoot can be killed, causing a "dead heart". In older plants the upper part of the stem usually dies as a result of the boring of the caterpillars.

Egg: The eggs are laid on the underside of a leaf near the midrib in 3-5 rows, in groups of 50-100. They are flattened, ovoid, and about 0.8 mm long. Hatching takes place after 7-10 days.

Larva: The young caterpillars produce characteristic leaf windowing. Sometimes, the early stages mine in the leaves, causing yellow streaks. After a few days they bore down inside the funnel. They also may move down outside the stem and then bore into it just above an internode. In older plants the caterpillars sometimes live in the developed heads. They are creamy pink with groups of dark spots along the back. The head capsule is brown. When mature they are about 25 mm long. The larval period takes 28-35 days.

Pupa: Pupation takes place in a small chamber in the stem. The pupal period takes 7-10 days.

Adult: Adult moths have a wingspan of 20-30 mm. Males are smaller and darker than females. The forewings of males are pale brown. The forewings of the females are much paler and the hind wings are almost white.

ETL for *Chilo partellus* is 6 – 8 No's of moths per trap per day.

Use Phormone Traps from 1 month crop stage @ 8 No's per acre to control pest at early stage. Trap canopy should be placed one foot above crop canopy to achieve optimum catch.

Plot No.23, TSIIC Techpark, Road No.15, IDA Nacharam, Hyderabad – 500076, Telangana, India

Tel: +91-40-27177918, Cell: +91-94408 97918

E-mail: info@pheromonechemicals.in, Web: www.pheromonechemicals.in

e-shop: www.pheromonechemicals.org



Pheromone Chemicals

The name you can always trust

Mfrs: Pheromone Traps, Lures, Yellow sticky traps

Lure Specifications:

1. Lures made of Virgin Silicone rubber for uniform release and long life
2. Minimum pheromone loading assured is 3 mg per lure
3. Shelf life of 18 months from date of manufacturing date at room temperature
4. Field efficacy will be 30-45 days after installation in field
5. Packed in trilaminated aluminum foil (LD, Aluminum, Polyester)
6. Lures will attract target pest species only

Use 8 No's of traps per acre.

Always use Phero – Sensor™ – SP / BP for best results

PC Pheromone Lures for
CHILO PARTELLUS
(Spotted Stalk borer)
Host – Sorghum, Maize, millet, Sugarcane, Rice

The advertisement contains two photographs: the left one shows a light-colored moth with its wings spread on a green leaf, and the right one shows a dark, segmented larva on a plant stem.

Plot No.23, TSIIC Techpark, Road No.15, IDA Nacharam, Hyderabad – 500076, Telangana, India

Tel: +91-40-27177918, Cell: +91-94408 97918

E-mail: info@pheromonechemicals.in, Web: www.pheromonechemicals.in

e-shop: www.pheromonechemicals.org